

FIG. 1 Ribbon representation of the structure of the murine antibody against canine lymphoma determined by X-ray analysis of the triclinic crystals. The heavy chains are shown in yellow and blue, the light chains in red. The stem of the molecule projects towards the viewer and assumes an asymmetric, oblique orientation with respect to the Fabs. This orientation illustrates the large difference in hinge angles of about 65° and 115° . The local dyad relating the heavy chains of the Fc is that dyad indicated by the primary solution of the self-rotation function. Fab2 is viewed along the axis through the switch peptides. Fab2 has an elbow angle of 143° , in contrast to Fab1 which has an elbow of 159° . Twenty-three residues in each heavy chain comprising the hinge regions seen here, were built into the model with idealized geometry using both FRODO²⁸ and XPLOR^{14,15}. These residues were missing from the fragment models taken from the Brookhaven Data Bank.

FIG. 2 *a*, stereo diagram of the monoclonal antibody viewed perpendicular to the approximate 2-fold axis relating the constant domains of the Fabs. This dyad was that indicated by the secondary solution of the self-rotation function. Apparent here is the difference in the two elbow angles and the consequent failure of the variable domains to maintain this relationship. Also apparent in this view is the failure of the Fc dyad to intersect the 2-fold axis relating the constant domains of the Fabs. Both symmetry axes are apparently independent local dyads. *b*, Stereo diagram of the IgG2a antibody showing the region between the CH2 domains. In the human IgG1 Fc fragment⁶, carbohydrate was located in this area between the two CH2 domains and is probably in a similar location in this antibody. No attempt has yet been made to include the carbohydrate component in the model. It can also be seen here that the dyad axis of the Fc does not intersect the approximate long axis of the Fabs. Colour coding is the same as in Fig. 1.




TABLE 1 Crystallographic data, structure solution and refinement

Crystal data: Space group $P1$; $a=66.39$ Å, $b=77.34$ Å, $c=101.42$ Å, $\alpha=87.6^\circ$, $\beta=92.6^\circ$, $\gamma=97.5^\circ$, $Z=1$; resolution ~ 3.0 Å. Data collection: SDMS (Xuong-Hamlin) detectors, Rigaku Ru-200 source, frame size 0.12° , counting time 60–120 s. Total observations, 114,867; at 3.5 Å unique reflections, 24,808; 98.7% complete; $R_{\text{sym}}=0.098$. After $F/\sigma=4.0$ cutoff, unique reflections at 3.5 Å = 20,964

Operation	Result
Self-rotation function	Two consistent dyad solutions in several resolution ranges
Fc rotation function	
(1) Search probe entire Fc	Two peaks related by pseudodyad; r.m.s.* = 6.03 and 5.54
(2) C α 3 domains	same solutions as (1); r.m.s. = 5.66 and 5.44
(3) C α 2 domains	same solutions as (1); r.m.s. = 3.41 and 3.14
Fab1 rotation function	
(1) Search probes were constant domains of 7 different Fabs	Exceptional peak with constant domains of HYHEL-5; r.m.s. = 6.42
(2) Probes were variable domains of 7 different Fabs	Solution optimal for variable domains of McPC603; r.m.s. = 4.35
(3) An intact Fab1 constructed according to above results	Outstanding solution consistent with (1) and (2); r.m.s. = 7.98
Fab2 rotation function	
Range of Fab models with elbows of 120° – 180° constructed using XPLOR by altering Fab1 elbow every 5°	Unambiguous solution for probe with elbow angle 140° ; r.m.s. = 6.27
Translation function	
(1) Fc fixed, Fab1 moving	A self-consistent set of solutions
(2) Fc fixed, Fab2 moving	from all searches (1)–(4);
(3) Fab1 fixed, Fab2 moving	$cc^\dagger=0.157$ – 0.249 for 4–8 Å
(4) Fab1 and Fab2 fixed, Fc moving	resolution
Refinement	
(1) Rigid body with twelve β -barrel domains; 3.5–12 Å	$R=0.386$ $cc^\dagger=0.529$
(2) Powell minimization and simulated annealing after insertion of correct amino-acid sequence (occupancy of 46 hinge residues set to zero); 3.5–8 Å	$R=0.188$ $cc^\dagger=0.876$
	r.m.s. deviations
	Bonds 0.017 Å
	Angles 4.038°
	Dihedrals 28.597°
	Impropers 0.686°

*All r.m.s. values are stated for 4–8 Å resolution searches. † cc, Correlation coefficient.


solutions were frequently found even with the probes representing one sixth of the antibody. Rotation function solutions were based on the human Fc fragment⁴, the constant domains of Fab HYHEL-5 (ref. 18), and the variable domains of Fab McPC603 (ref. 19), including the hypervariable regions. Translation searches were performed to determine the relative distances between the three portions of the molecule, the two Fabs and the Fc (Table 1).

The antibody structure was assembled according to essentially independent, but internally consistent, molecular replacement results that ultimately yielded a model fully consistent with the

stereochemistry of an intact antibody. The packing revealed good complementarity of surfaces without interpenetration. Lattice contacts immobilize all segments of the molecule to permit visualization of both Fabs as well as the Fc.

The structure of the antibody is shown in Figs 1 and 2. Its most prominent features are: (1) There is an approximate 2-fold axis relating the heavy chains of the Fc portion of the molecule. The dyad deviates particularly for the CH2 domains; (2) the disposition of the Fc with respect to the Fab portions is quite oblique; (3) the hinge angle between the Fc and Fab1 is approximately 65° and for Fab2 about 115° ; (4) the long axes of the

FIG. 3 Stereo diagram of the packing of four antibodies in the triclinic cell, each of a different colour, showing the intricate network of intermolecular contacts that stabilizes the conformation of the molecule. The Fc segments, which lie more or less along the longest body diagonal, are immobilized by multiple Fab contacts, suggesting why the Fc in these crystals is ordered. The constant domains of an Fab2 of one molecule insert in the elbow region of Fab1 of a different antibody molecule to fix the dispositions of the Fabs. Notable initial exceptions to the otherwise acceptable packing were three hypervariable loops protruding from the variable domains of Fab McPC603. When the correct sequence for the canine lymphoma antibody was examined, it was apparent that the offending residues corresponded to deletions in the latter molecule. Thus, when the correct amino-acid sequence was substituted, virtually all of the packing exceptions were eliminated, as shown in this view.



two Fabs are almost collinear; thus, there is an approximate long axis running through the entire Fab assembly. The angle between the Fabs is $170 \pm 2^\circ$, and the Fab axes are offset by 9 Å; (5) Fab1 has an elbow angle of 159° , and Fab2 has an elbow angle of 143° . These elbow angles are near the middle of the range of values observed for other Fabs¹; (6) the constant domains of Fab1 and Fab2 are related by a near exact dyad axis of symmetry. The variable domains are not so related because of the difference in elbow angles of the Fabs; (7) the dyad of the Fc is at an angle of about 120° with that dyad relating constant Fab domains; (8) the Fc 2-fold axis does not intersect the dyad relating the constant domains of the Fabs, nor does it intersect the approximate long axis of the Fabs; (9) in the crystal, all segments share extensive interfaces which severely restrict the dispositions of neighbours. The contacts, illustrated in Fig. 3, presumably stabilize this particular conformation.

The asymmetric conformation, observed in these crystals of the antitumour antibody, should probably not be considered as a static structure which is maintained in solution. The structure probably represents only one of many possible transient conformations. The unique structure is a product of the intrinsic flexibility of the antibody and the lattice interactions that stabilize this particular distribution of domains. Indeed, electron microscopy²⁰⁻²³, fluorescence polarization^{24,25} and previous X-ray crystallographic studies^{5,26,27} have provided extensive evidence for a wide range of conformations based on segmental flexibility.

The structure we present is instructive in that it illustrates the nature and extent of this structural variability, or dynamic range, which is inherent in the antibody. The Fabs are loosely tethered to a mobile Fc. Each Fab can assume its own elbow angle as its environment or function requires. Somewhat unexpected is the fact that, were the elbow angles the same, the Fabs would be related by an almost exact 2-fold axis that is quite independent of the Fc. It is the disposition of the Fc that disrupts the overall symmetry of the molecule. This is in keeping with the disorder, or multiple orientations of the Fc, observed in the Kol antibody structure^{7,8}.

The hinge polypeptides are not really hinges, but rather they are tethers that allow the Fab components to drift from the Fc to bind antigen or potentially allow the Fc to move in such a way to trigger effector functions, such as the activation of complement^{25,28}. The connecting polypeptides give the Fabs the freedom to move and twist so as to align hypervariable regions with antigenic sites on large, immobile carriers, in this case tumour cells. The crystal structure visually demonstrates that the antibody is an assembly of units possessing a high degree of flexibility, a molecule suited to the task of scavenging foreign objects or activating a cell lysis system. □

Received 3 August; accepted 9 October 1992.

- Wilson, L. A., Rini, J. M., Fremont, D. H., Flesher, G. G. & Stura, E. A. *Meth. Enzym.* **203**, 153-176 (1991).
- Alzari, P. M., Lascombe, M.-B. & Poljak, R. J. *A. Rev. Biochem.* **6**, 555-580 (1988).
- Davies, D. R., Padlan, E. A. & Sheriff, S. A. *Rev. Biochem.* **59**, 439-473 (1990).
- Deisenhofer, J. *Biochemistry* **20**, 2361-2370 (1981).
- Silverton, E. W., Navia, M. A. & Davies, D. R. *Proc. natn. Acad. Sci. U.S.A.* **74**, 5140-5144 (1977).
- Rajan, S. S. et al. *Mol. Immun.* **20**, 797-799 (1983).
- Colman, P. M., Deisenhofer, J., Huber, R. & Palm, W. *J. molec. Biol.* **100**, 257-282 (1976).
- Marquardt, M., Deisenhofer, J., Huber, R. & Palm, W. *J. molec. Biol.* **141**, 389-391 (1980).
- Steplewski, Z., Jeglum, K. A., Rosales, C. & Weintraub, N. *Cancer Immun. Immunother.* **24**, 197-201 (1987).
- Rosales, C., Jeglum, K. A., Obrodka, M. & Steplewski, Z. *Cell. Immun.* **115**, 420-428 (1988).
- Jeglum, K. A. *Proc. Seventh AIVM Forum*, San Diego (922-925) (Lippincott, Hagerstown, Maryland, 1989).
- Larson, S., Day, J., Greenwood, A., Skaletsky, E. & McPherson, A. *J. molec. Biol.* **222**, 17-19 (1991).
- Fitzgerald, P. M. D. *J. appl. Crystallogr.* **21**, 273-276 (1988).
- Brunker, A. T., Kurlyan, J. & Karplus, M. *Science* **235**, 458-460 (1987).
- Brunker, A. T. *A. Rev. phys. Chem.* **42**, 197-223 (1991).
- Bernstein, F. C. et al. *J. molec. Biol.* **112**, 535-542 (1977).
- Crowther, R. A. in *The Molecular Replacement Method* (ed. Rossmann, M. G.) 173-178 (Gordon and Breach, New York, 1972).
- Sheriff, S. et al. *Proc. natn. Acad. Sci. U.S.A.* **84**, 8075-8079 (1987).
- Satow, Y., Cohen, G. H., Padlan, E. A. & Davies, D. R. *J. molec. Biol.* **190**, 593-604 (1986).
- Wrigley, N. G., Brown, E. B. & Skehel, J. J. *J. molec. Biol.* **169**, 771-774 (1983).
- Roux, K. H. *Eur. J. Immun.* **14**, 459-464 (1984).
- Lamy, J. et al. *Biochemistry* **24**, 5532-5542 (1985).

- Wade, R. H., Taveau, J. C. & Lamy, J. N. *J. molec. Biol.* **206**, 349-356 (1989).
- Schneider, W. P., Wensel, T. G., Stryer, L. & Oi, V. T. *Proc. natn. Acad. Sci. U.S.A.* **85**, 2509-2511 (1988).
- Dangl, J. L. et al. *EMBO J.* **7**, 1989-1994 (1988).
- Huber, R., Deisenhofer, J., Colman, P. M. & Matsushima, M. *Nature* **264**, 415-420 (1978).
- Amzel, L. M. & Poljak, R. J. *A. Rev. Biochem.* **48**, 951-997 (1979).
- Oi, V. T. et al. *Nature* **307**, 136-140 (1984).
- Jones, T. A. in *Computational Crystallography* (ed. Sayre, D.) 307-317 (Oxford University Press, New York, 1982).

ACKNOWLEDGEMENTS. This research was supported by grants from the NIH from the NSF and NASA, as well as by the Immunopharmaceuticals Company of San Diego. We thank the San Diego Supercomputer Facility for time on the Cray Y-MP.

RETRACTION

Identification by anti-idiotypic antibodies of an intracellular membrane protein that recognizes a mammalian endoplasmic reticulum retention signal

D. Vaux, J. Tooze & S. Fuller

Nature **345**, 495-502 (1990)

OUR further characterization of the *M*_r 72,000 (72K) protein has shown that the data in Fig. 2 of our paper are erroneous and not repeatable. We retract the statement that the 72K protein is an integral membrane protein. The present evidence is consistent with this protein being associated with the intermediate compartment. We also withdraw our speculation concerning its function.

GUIDE TO AUTHORS

PLEASE follow these guidelines to ensure your manuscript can be handled expeditiously. *Nature* is an international journal covering all the sciences. Contributions should therefore bear in mind those readers who work in other fields and those for whom English is a second language, and write clearly and simply, avoiding unnecessary technical terminology. Space in the journal is limited, making competition for publication severe. Briefly a highly valued one printed page of *Nature*, without interruptions, contains about 1,300 words.

Manuscripts are selected for publication according to editorial assessments of their scientific and originality. Reviews are invited. They can be sent to London or Washington and should be addressed to the Editor. Manuscripts may be dealt with in either office, depending on the subject matter, and will where necessary be sent between offices by overnight courier. High priority cannot be given to prepublication enquiries; in urgent cases they can be made in the form of a one-page fax. All manuscripts are acknowledged on receipt but fewer than half are sent for review. Those that are not reviewed are returned as rapidly as possible so that they may be submitted elsewhere without delay. Contributors may suggest reviewers; limited requests for the evaluation of specific reviewers are usually heeded. Manuscripts are usually sent to two or three reviewers, chosen for their expertise rather than their geographical location. Manuscripts accepted for publication are typeset from the London office.

Nature requests authors to deposit sequence and crystallographic data in the database that exist for this purpose, and to mention availability of these data.

Once a manuscript is accepted for publication, contributors will receive proofs in about 4 weeks. *Nature's* staff will edit manuscripts with a view to brevity and clarity; contributions should check proofs carefully. Manuscripts are generally published 2-3 weeks after receipt of corrected proofs. *Nature* does not read page charges. Contributors receive a reprint order form with their proofs; reprint orders are processed after the manuscript is published and payment received.

Categories of paper

Review articles survey recent developments in a field. Most are commissioned and originate as editorials. Some are invited. They are synopsis addressed to the Reviews Coordinator. Length is negotiable to advance.

Articles are research reports whose conclusions are of general interest and which are sufficiently rounded to be a substantial advance in understanding. They should not have more than 3,000 words of text (not including figure legends) or more than six display items; they should not occupy more than five pages of *Nature*.

Articles start with a heading of 30-50 words written to advertise their content in general terms, to which editors will pay particular attention. The heading does not usually contain numbers, abbreviations or measurements. The introduction to the study is contained in the first two or three paragraphs of the article, which also briefly summarizes its results and implications. Articles have fewer than 30 references and may contain a few short subheadings.

Letters are short reports of outstanding novel findings whose implications are general and important enough to be of interest to those outside the field. Letters should have 1,000 or fewer words of text and four or fewer display items. The first paragraph describes, in not more than 150 words, the nature of the observations, the background, rationale and chief conclusions of the study for the particular benefit of non-specialist readers. Letters do not have subheadings and should contain fewer than 30 references.

Commentary articles deal with issues in, or arising from, research that are of interest to readers. They are invited and are distinguished by their suggestions can be made to the Commentary Editor in the form of a one-page synopsis.

News and View articles inform non-specialist readers of scientific advances, sometimes in the form of a column. News items are commissioned but proposals can be made to editors. News and View Editor.

Scientific Correspondence is for discussion of topical scientific issues. Priority is given to letters of fewer than 300 words.

Preparation of manuscripts

All manuscripts should be typed, double-spaced, on one side of paper only. An original and four copies are required, all accompanied by artwork. If photographs are included, five or eight are required; for line drawings, one set of originals and five or eight glossy photographs are acceptable. References to figures should be provided. Suggestions for cover illustrations, if any, should be provided. All figures should be on separate sheets, all of which should be typed and numbered. Three copies of microfilm may be submitted for publication elsewhere should be included with a copy of a submitted manuscript, and clearly marked as such. Copies of revised and resubmitted manuscripts, typed with manuscript numbers are required, together with legends and detailing the changes made.

Titles are brief and simple. Active verbs, nominalized verbs, adjectives and punctuation are to be avoided. They should consist of two or three key words for indexing purposes.

Abstracts should be marked individually and clearly with their nature and, where known, the manuscript number. They should be larger than 25 by 25 cm. Figures with small parts should be avoided and are permitted only if the parts are directly related experimentally or logically. Unrelated original images should be provided. Suggestions for cover illustrations, if any, should be provided with the manuscript, together with legends and detailing the changes made.

Protein/nucleotide sequences should ideally be in the text and not the single-letter code for amino acids. One alternative is to use a separate sheet for each sequence, typed and numbered.

Colour artwork. A charge of £200 per page is made at the time of the proof to cover the cost of reproducing colour figures. Facilities to reduce costs will not prevent publication of essential colour figures if circumstances are explained. Proofs of colour artwork should be submitted with the manuscript, together with legends and detailing the changes made. **Figure legends** should not exceed 300 words and should ideally be shorter. The figure is described first, then, briefly, legend text is given. References are numbered sequentially as they appear in the text, followed by numbers in tables and finally by those in figures in papers published first or in the press are numbered and listed in the reference list. All other forms of reference should be listed in the text as a personal communication, manuscript submitted or in press. Text is not included in reference lists. References are given according to the *World List of Scientific Periodicals* (London, 1963-65). The first and last page numbers of each reference to books should include publisher, place and year.

Abbreviations, symbols, units and Greek letters should be used in the first time they are used. Acronyms should be used wherever possible and, if used, defined. Footnotes are not used in *Nature*.

Acknowledgements are brief and appear after the references and contribution numbers are not allowed. **Supplementary information** is material relevant to the article, which cannot, for lack of space, be published in full, but is available from *Nature* on request.

Submission. Manuscripts can be sent to the Editor at a Life Sciences, London WC2N 3LP, UK or at 1224 National Press Building, Washington, DC 20045, USA. Manuscripts or proofs sent by courier to London should be declared in manuscript and value to prevent the imposition of import duty and value-added tax.